arXiv:1112.4315v1 [math.CT] 19 Dec 2011

A characterization of the category Q-**TOP**

Sheo Kumar Singh*
Department of Mathematics,
Banaras Hindu University,
Varanasi-221005, India

Arun K. Srivastava†
Department of Mathematics
and
Centre for Interdisciplinary Mathematical Sciences,
Banaras Hindu University,
Varanasi-221005, India

1 Introduction

E.G. Manes in [2] gave a somewhat 'axiomatic' characterization (upto isomorphism) of the category **TOP** of topological spaces, among a certain class of categories, satisfying some conditions, in which a Sierpinski space-like object, played a key role. Following that, the category $[0,1]$-**TOP** of $[0,1]$-topological spaces (known more commonly as 'fuzzy' topological spaces) was analogously characterized by Srivastava [4]. S.A. Solovyov [3] has recently introduced the notion of Q-topological spaces (Q being a fixed member of a variety of Ω-algebras). In this note, we give a characterization of the category Q-**TOP** of Q-topological spaces in which the so-called Q-Sierpinski space, introduced in [3], plays a key role.

2 The category Q-TOP

We begin by recalling the (well-known) notions of Ω-algebras and their homomorphisms. For details, see [1].

*sheomathbhu@gmail.com
†arunksrivastava@gmail.com

Definition 2.1 ([3])

- *Let $\Omega = (n_\lambda)_{\lambda \in I}$ be a class of cardinal numbers. An* **Ω-algebra** *is a pair $(A, (\omega_\lambda^A)_{\lambda \in I})$ consisting of a set A and a family of maps $\omega_\lambda^A : A^{n_\lambda} \to A$. Any $B(\neq \phi) \subseteq A$ is called a* **subalgebra** *of $(A, (\omega_\lambda^A)_{\lambda \in I})$ if for any $\lambda \in I$ and any $(b_i)_{i \in n_\lambda} \in B^{n_\lambda}$, $\omega_\lambda^A((b_i)_{i \in n_\lambda}) \in B$. Given $S \subseteq A$, the intersection of all the subalgebras of $(A, (\omega_\lambda^A)_{\lambda \in I})$ containing S is clearly a subalgebra of $(A, (\omega_\lambda^A)_{\lambda \in I})$. We shall denote it as $< S >$.*

- *Given $(A, (\omega_\lambda^A)_{\lambda \in I})$ and $(B, (\omega_\lambda^B)_{\lambda \in I})$, a map $f : A \to B$ is called an* **Ω-algebra homomorphism** *if for every $\lambda \in I$ the following diagram commutes:*

$$\begin{array}{ccc} A^{n_\lambda} & \xrightarrow{f^{n_\lambda}} & B^{n_\lambda} \\ {\scriptstyle \omega_\lambda^A}\downarrow & & \downarrow{\scriptstyle \omega_\lambda^B} \\ A & \xrightarrow[f]{} & B \end{array}$$

 .
 Let **Alg**(Ω) *denote the category of Ω-algebras and Ω-algebra homomorphisms.*

- *A* **variety** *of Ω-algebras is a full subcategory of* **Alg**(Ω) *which is closed under the formation of products* [1]*, subalgebras and homomorphic images.*

Throughout, Q denotes a fixed member of a fixed variety of Ω-algebras.

- *Given a set X, a subset τ of Q^X is called a* **Q-topology** *on X if τ is a subalgebra of Q^X; in such a case, the pair (X, τ) is called a* **Q-topological space**.

- *Given two Q-topological spaces (X, τ) and (Y, η), a* **Q-continuous function** *from (X, τ) to (Y, η) is a function $f : X \to Y$ such that $f^{\leftarrow}(\alpha) \in \tau, \forall \alpha \in \eta$, where $f^{\leftarrow}(\alpha) = \alpha \circ f$.*

It is evident that all Q-topological spaces, together with Q-continuous maps, form a category, which we shall denote as Q-**TOP**.

The following categorical concepts are from Manes [2]

Definition 2.2 ([2])

1. *A* **category C of sets with structure** *is defined through the following descriptions of its objects ('C-structured sets') and morphisms ('C-admissible maps'), satisfying the two axioms given below:*
 A class $C(X)$ of 'C-structures' is assigned with each set X and a 'C-structured set' is a pair (X, s), with $s \in C(X)$.

[1] The category **Alg**(Ω) is closed under products

A subset $C(s,t)$ of the set of all functions from X to Y is assigned with each pair of C-structured sets (X,s) and (Y,t) and a 'C-admissible map' from (X,s) to (Y,t) is any $f \in C(s,t)$ (in which case we write "$f : (X,s) \to (Y,t)$").
The axioms are:
Axiom A_1*: If $f : (X,s) \to (Y,t)$ and $g : (Y,t) \to (Z,u)$ then also $g \circ f : (X,s) \to (Z,u)$.*
Axiom A_2*: Given a bijection $f : X \to Y$ and $t \in C(Y)$, there exists a unique $s \in C(X)$ such that $f : (X,s) \to (Y,t)$ and $f^{-1} : (Y,t) \to (X,s)$.*

2. *Given a category* **C** *of structures (as defined above), a family $\mathcal{F} = \{f_j : (X,s) \to (Y_j,t_j) | j \in J\}$ of C-admissible maps is said to be* **optimal** *if for each C-structured set (Z,u) and a function $g : Z \to X$, $g : (Z,u) \to (X,s)$ iff $f_j \circ g : (Z,u) \to (X,s), \forall j \in J$. Further, if for a family $\mathcal{F} = \{f_j : X \to (Y_j,t_j) | j \in J\}$ of functions, where X is a set and each (Y_j,t_j) is a C-structured set, there exists $s \in C(X)$ such that the family $\{f_j : (X,s) \to (Y_j,t_j) | j \in J\}$ is optimal, then s is called an* **optimal lift** *of the family $\mathcal{F}$.*

3. *An object $S = (S,u)$ in a category C of sets with structure is called a* **Sierpinski object** *if for every C-object $X = (X,s)$, the family of all C-admissible maps from X to S is optimal.*

Remark 2.1 *It is easy to verify that, in Q-***TOP***, the optimal lift of a family $\mathcal{F} = \{f_j : X \to (Y_j,t_j) | j \in J\}$ of functions, where X is a set and each (Y_j,t_j) is a Q-***TOP***-object, is precisely the smallest Q-topology on X making each f_i Q-continuous.*

Both **TOP** and $[\mathbf{0},\mathbf{1}]$-**TOP** are categories of sets with structures and the usual two-point Sierpinski space and the fuzzy Sierpinski space (of [4]) are Sierpinski objects in these categories.

We verify (on expected lines) that Q-**TOP** is also a category of sets with structures. For each set X, $C(X)$ is the family of all Q-topologies on X and the C-admissible maps are just the Q-continuous maps. Given a bijection $f : X \to Y$ and $t \in C(Y)$, there exists a unique $s \in C(X)$, namely $s =< \mathscr{A} >$, with $\mathscr{A} = \{q \circ f | q \in t\}$, such that both $f : (X,s) \to (Y,t)$ and $f^{-1} : (Y,t) \to (X,s)$ are Q-continuous. Also, every family $\mathcal{F} = \{f_j : X \to (Y_j,t_j) | j \in J\}$ of functions, where X is a set and (Y_j,t_j), $j \in J$ are the Q-topological spaces, has an optimal lift s, namely $s =< \mathscr{S} >$, with $\mathscr{S} = \{q \circ f_j | q \in t_j, j \in J\}$. The *product* of a family $\{(X_i,t_i) | i \in I\}$ of Q-topological spaces is the Q-topological space $(\prod X_i, t)$, where t is the optimal lift of the family of all the projection maps $p_i : \prod X_i \to (X_i,t_i)$. In fact, $(\prod X_i, t)$ is the categorical product of the Q-**TOP**-objects (X_i,t_i), $i \in I$.

3 Q-Sierpinski Space

Let $S = Q$. It is obvious that $u =< id >$, where $id \in Q^Q$ is the identity function, is a Q-topology on S. (S, u) has been called the Q-**Sierpinski space** in [3]. Theorem 3.2 establishes the appropriateness of this concept of Q-Sierpinski space. But first, we state the following easy-to-verify result.

Theorem 3.1 *For any Q-topological space (X, τ), $p \in \tau$ iff $p : (X, \tau) \to (S, u)$ is Q-continuous.*

Theorem 3.2 *(S, u) is a Sierpinski object in Q-**TOP**.*

Proof: Let (X, τ) be a Q-**TOP**-object and $\mathcal{F} = \{f : (X, \tau) \to (S, u) | f$ is Q-continuous$\}$ be the family of all Q-continuous maps from (X, τ) to (S, u). To show that this family is optimal, let τ' be any other Q-topology on X making each $f \in \mathcal{F}$ Q-continuous. If $p \in \tau$ then, as $id \in u$, $p : (X, \tau) \to (S, u)$ must be Q-continuous and so $p : (X, \tau') \to (S, u)$ is also Q-continuous. But then $p \in \tau'$. Thus $\tau \subseteq \tau'$, showing that τ is the smallest Q-topology on X making each $f \in \mathcal{F}$ Q-continuous. □

4 A characterization of Q-TOP

Theorem 4.1 *A category* **C** *of sets with structures is isomorphic to the category Q-**TOP** if and only if* **C** *contains an object (S, u), with $S = Q$, satisfying the following conditions:*

1. *every family $f_i : X \longrightarrow (S, u)$ has an optimal lift,*
2. *the maps $\omega_\lambda : (S, u)^{n_\lambda} \longrightarrow (S, u)$ are* **C***-admissible for each $\lambda \in I$, where $(S, u)^{n_\lambda} = (S^{n_\lambda}, u_{n_\lambda})$, with u_{n_λ} being the optimal lift of all the projection maps from S^{n_λ} to (S, u),*
3. *(S, u) is a Sierpinski object in* **C**.

Proof: First, it is clear that Q-**TOP** satisfies (1) and (3). That Q-**TOP** also satisfies (2) is shown as follows. Note that u_{n_λ}, the product Q-topology on S^{n_λ} has a generating set consisting precisely of all the projection maps from S^{n_λ} to S. To show that $\omega_\lambda : (S, u)^{n_\lambda} \longrightarrow (S, u)$ is Q-continuous, it suffices to show that $\omega_\lambda \in u_{n_\lambda}$. Now if $a = (a_i)_{i \in n_\lambda} \in S^{n_\lambda}$, then ${\omega_\lambda}^{S^{S^{n_\lambda}}}((p_i)_{i \in n_\lambda})(a) = \omega_\lambda^S((p_i(a))_{i \in n_\lambda}) = \omega_\lambda^S((a_i)_{i \in n_\lambda}) = \omega_\lambda(a)$, where ${p_i}'s$ are the projection maps from S^{n_λ} to S. Hence $\omega_\lambda = {\omega_\lambda}^{S^{S^{n_\lambda}}}((p_i)_{i \in n_\lambda}) \in u_{n_\lambda}$. Thus ω_λ is Q-continuous, $\forall \lambda \in I$.

For each set X, let $Q(X)$ denote the set of all Q-topologies on X. It will suffice to prove that, for each set X, there exists a bijection $\Phi_X : C(X) \to Q(X)$ such that $f : (X, s) \to (Y, t)$ is C-admissible iff $f : (X, \Phi_X(s)) \to (Y, \Phi_Y(t))$ is Q-continuous, i.e., iff $q \circ f \in \Phi_X(s)$ for each $q \in \Phi_Y(t)$. For each set X and

$s \in C(X)$, put $\Phi_X(s) = \{p \in Q^X | p : (X, s) \to (S, u)$ is C-admissible$\}$. We show that $\Phi_X(s)$ is a Q-topology on X, i.e., $\Phi_X(s)$ is a subalgebra of Q^X, or that $\Phi_X(s) \in Q(X)$. For a given $\lambda \in I$, consider $(p_i)_{i \in n_\lambda} \in (Q^X)^{n_\lambda}$, with $p_i \in \Phi_X(s)$, $\forall i \in n_\lambda$. We wish to show that $\omega_\lambda^{Q^X}((p_i)_{i \in n_\lambda}) \in \Phi_X(s)$, i.e., $\omega_\lambda^{Q^X}((p_i)_{i \in n_\lambda})$ is also a C-admissible map from (X, s) to (S, u). The map $f : (X, s) \to (S, u)^{n_\lambda}$, defined by $f(x)(j) = p_j(x)$, for each $x \in X$ and $1 \leq j \leq n_\lambda$, is easily seen to be C-admissible. Here we note that $\omega_\lambda^{Q^X}((p_i)_{i \in n_\lambda}) = \omega_\lambda \circ f$, as $(\omega_\lambda \circ f)(x) = \omega_\lambda(f(x)) = \omega_\lambda^Q((f(x)(i))_{i \in n_\lambda}) = \omega_\lambda^Q((p_i(x))_{i \in n_\lambda}) = \omega_\lambda^{Q^X}((p_i)_{i \in n_\lambda})(x)$. So, $\omega_\lambda^{Q^X}((p_i)_{i \in n_\lambda})$ is also a C-admissible, being the composition of two C-admissible maps. Thus $\Phi_X(s)$ is a Q-topology on X.

Let $\tau \in Q(X)$, $\mathcal{F} = \{f : (X, \tau) \to (S, u) | f$ is Q-continuous$\}$, and s_τ be the optimal lift of the family $\{f : X \to (S, u) | f \in \mathcal{F}\}$. Then clearly, $s_\tau \in C(X)$ and $f : (X, s_\tau) \to (S, u)$ is C-admissible for each $f \in \mathcal{F}$. This provides a function from $Q(X)$ to $C(X)$, which, it can be verified, is the inverse of $\Phi_X : C(X) \to Q(X)$. So, $C(X)$ and $Q(X)$ are in one-to-one correspondence. Also, condition (3) says that $f : (X, s) \to (Y, t)$ is C-admissible precisely when $q \circ f \in \Phi_X(s)$ for each $q \in \Phi_Y(t)$.

References

[1] N. Jacobson, Basic Algebra-II, W.H. Freeman and Company, 1980.

[2] E.G. Manes, "Algebraic Theories", Springer-Verlag, New York, 1976.

[3] S.A. Solovyov, Sobriety and spatiality in the variety of algebras, *Fuzzy Sets and System*, **159** (2008), 2567-2585.

[4] A.K. Srivastava, Fuzzy Sierpinski space, *J. of Math. Anal. and Appl.*, **103** (1984), 103-105.